TABLE 6a. 2004 Reported Deer Kill by CDF&G Region, Sex, Antler Class and Percent by Antler Class of Bucks (a).

		Unclassified		Spike		2 Point		3 Point		4 Point		> 4 Point			
		Point C	oint Count (b)		Bucks		Bucks		Bucks		Bucks		Bucks		
CDF&G	Total		% of		% of		% of		% of		% of		% of	Total	TOTAL
Region	Region	Region	Region	Region	Region	Region	Region	Region	Region	Region	Region	Region	Region	Region	REGION
of	Doe	Buck	Buck	Buck	Buck	Buck	Buck	Buck	Buck	Buck	Buck	Buck	Buck	Buck	DEER
Kill	Kill	Kill	Kill	Kill	Kill	Kill	Kill	Kill	Kill	Kill	Kill	Kill	Kill	Kill	KILL
Region 1 - Northern California and North Coast	10	47	0.7%	3	0.0%	3,066	42.8%	2,474	34.5%	1,376	19.2%	195	2.7%	7,161	7,171
Region 2 - Sacramento Valley and Central Sierra	52	40	1.1%	6	0.2%	1,790	48.9%	1,087	29.7%	622	17.0%	119	3.2%	3,664	3,716
Region 3 - Central Coast	47	14	0.3%	11	0.3%	2,614	60.5%	1,210	28.0%	424	9.8%	46	1.1%	4,319	4,366
Region 4 - San Joaquin Valley and Southern Sierra	0	22	0.8%	1	0.0%	1,445	50.6%	810	28.4%	467	16.3%	112	3.9%	2,857	2,857
Region 5 - South Coast	182	22	2.2%	3	0.3%	658	66.2%	226	22.7%	73	7.3%	12	1.2%	994	1,176
Region 6 - Eastern Sierra and Inland Deserts	5	3	0.3%	3	0.3%	440	45.5%	264	27.3%	209	21.6%	48	5.0%	967	972
STATEWIDE TOTALS:	296	148	0.7%	27	0.1%	10,013	50.2%	6,071	30.4%	3,171	15.9%	532	2.7%	19,962	20,258

TABLE 6a NOTES:

2004table6a.xls Page 1 of 1

⁽a) Table 6a does not include PLM kill data. PLM kill analysis is included in TABLES 7a through 7f.

⁽b) Unclassified Point Count may be bucks without a reported point count or bucks with no points (antlerless and either-sex hunts).